

Preliminary results on the reaction of the ${}^6\text{He}+p$ scattering experiment in MS

The Machine study for the ${}^6\text{He}+p$ experiment was carried out in CRIB, the ${}^6\text{He}$ beam energy was ~ 48 MeV with 91% purity via 20 μm mylar degrader. We also carried out one short scattering measurement of the ${}^6\text{He}+p$ reaction with one set of dE-E detectors, the locus of the p, d, t, alpha, ${}^6\text{He}$ can be clearly seen. The proton and triton were identified combined with the kinematic calculation of the ${}^6\text{He}(p,p){}^6\text{He}$ and ${}^6\text{He}(p,t){}^4\text{He}$ reactions. In addition, the cross section of the ${}^6\text{He}(p,p){}^6\text{He}$ and ${}^6\text{He}(p,t){}^4\text{He}$ reaction were roughly calculated, which is consistent with the tendency of the theoretical calculation.

Primary authors: ZHANG, Qian (Center for Nuclear Study, the University of Tokyo); YAMAGUCHI, Hidetoshi (Center for Nuclear Study, the University of Tokyo); SFERRAZZA, Michele (Department of Physics, Université Libre de Bruxelles); HAYAKAWA, Seiya (Center for Nuclear Study, the University of Tokyo); OKAWA, Kodai (Center for Nuclear Study, the University of Tokyo); CHERUBINI, Silvio (INFN - Laboratori Nazionali del Sud); LA COGNATA, Marco; PIZZONE, Gianluca; CHILLERY, Thomas William (Center for Nuclear Study, the University of Tokyo); HANAI, Shutaro (Center for Nuclear Study, the University of Tokyo); IMAI, Nobu (Center for Nuclear Study, the University of Tokyo); MASUOKA, Shoichiro (Center for Nuclear Study, the University of Tokyo); YAKO, Kentaro (Center for Nuclear Study, the University of Tokyo)

Track Classification: Experimental Nuclear Physics: Low and Intermediate Energies