

Mean-field study of the radiative capture $^{12}\text{C}(p,\gamma)^{13}\text{N}$ and $^{13}\text{C}(p,\gamma)^{14}\text{N}$ reactions

Friday, 24 August 2018 15:40 (15 minutes)

In this framework we study the effect of local optical potential on the radiative capture $^{12}\text{C}(p,\gamma)^{13}\text{N}$ and $^{13}\text{C}(p,\gamma)^{14}\text{N}$ reactions. The optical potential of nucleon-nucleus interaction is constructed by parameterization of Woods-Saxon potential and folding model using the effective nucleon-nucleon interaction CDM3Yn based on an extended Hartree-Fock calculation. The result indicates that the both potentials described effectively the (p,γ) reactions compared to the experimental data.

Primary author: Mr NGUYEN, Le-Anh (Ho Chi Minh City University of Education)

Co-authors: Mr NGUYEN, Tri-Toan-Phuc (Ho Chi Minh City University of Science); Mr NGUYEN, Hoang-Phuc (Institute for Nuclear Science and Technology, VINATOM); Ms DOAN, Thi-Loan (Institute for Nuclear Science and Technology, VINATOM); Prof. DAO, Tien-Khoa (Institute for Nuclear Science and Technology, VINATOM)

Presenter: Mr NGUYEN, Le-Anh (Ho Chi Minh City University of Education)

Session Classification: YSS