


# Single-particle and collective motions from nuclear many-body correlation (PCM2025)



Contribution ID: 78

Type: **not specified**

## Nuclear magnetic properties from first principles

*Tuesday, 4 March 2025 16:15 (25 minutes)*

Nuclear magnetic properties provide valuable insights into nuclear structure. In particular, the magnetic dipole moment is sensitive to how much the nucleus is dominated by the single-particle picture. Reproducing magnetic dipole moments has been one of the major challenges in nuclear *ab initio* theory. With the valence-space in-medium similarity renormalization group (VS-IMSRG), one of the *ab initio* calculation methods applicable for medium-mass and heavy nuclei, it was found that the absolute size of the magnetic dipole moments is underestimated. The effect of two-body current (2BC, also known as the meson exchange current) is non-negligible in light nuclei, as studied by Green's function Monte Carlo and no-core shell model. Thus, including 2BC effects in medium-mass and heavy nuclei calculations is a natural step forward. In this presentation, using the VS-IMSRG, I will discuss the 2BC effect on the magnetic dipole moments of nuclei near doubly magic systems from oxygen to bismuth. Additionally, I will present the effects of 2BC on magnetic transitions and form factors.

### Type of contribution

poster

### Are you a student or postdoc?

no

**Primary author:** MIYAGI, Takayuki (University of Tsukuba)

**Presenter:** MIYAGI, Takayuki (University of Tsukuba)

**Session Classification:** Session #4